

Epiphytic lichen diversity on *Populus tremula* L. (European aspen) in Uludağ mountain (Bursa)

Nevin HOCAOĞLU Şaban GÜVENÇ*

Uludağ University, Faculty of Arts and Sciences, Department of Biology, 16059 Nilüfer-Bursa / Turkey

*Corresponding author:
Email:saban@uludag.edu.tr

Received: December 25, 2015

Accepted: February 10, 2016

Abstract

A total of 40 epiphytic lichen species were found on *Populus tremula*. A total of 30 species were collected from the trunk base and 37 species from on the tree trunk at 150 cm above the ground. The most common species with the frequency of occurrence on the trees, respectively were *Lecidella elaeochroma*, *Melanohalea exasperatula*, *Caloplaca Cerina*, *Bacidia subincompta*, *Phlyctis argena*, *Physcia leptalea*, *Hypogymnia tubulosa* and *Lecanora hagenii*. The differences in the epiphytic lichen diversity between the base and trunk of *Populus tremula* is statistically significant, but is not statistically significant depending on the direction and circumference of tree trunk.

Keywords : Epiphytic lichens, diversity, community, Uludağ, *Populus tremula*

INTRODUCTION

As lichens are poikilohydric, they are not very efficient in controlling their water content or light capture efficiency and are, therefore, very sensitive to changes in microclimate [1, 2]. Recently, the epiphytic lichen communities has been found to be influenced by various factors as location, topography, climate, pollution, vegetation structure and composition [3, 4].

Epiphytic lichen diversity and species composition varies according to the structural diversity of the area, such as the different tree species in the same area, the age and viability of trees, cover, and microclimate [5, 6, 7]. The development of epiphytic lichens typically depends on the structural and chemical properties of the bark of trees on which they are grown [8, 9].

Lichen abundance in forests is controlled by environmental conditions, due to the influence of microhabitat conditions such as diameter, height and architecture [10]. Epiphytic lichen vegetation is strongly affected by altered microclimate, resulting from rapid changes in forest structure [11, 12]. The diversity of epiphytic species vary significantly, depending on the change of microclimatic parameters towards the inner part of the Mediterranean forest [13].

The aim of this study was to determine the differences in the species diversity and composition of the epiphytic lichens by the northern, southern, eastern and western direction on the base and trunk of *Populus tremula*. At the same time, variation in species diversity and composition were compared with the trunk base and trunk.

MATERIALS AND METHODS

Study area

Uludağ is the highest mountain in the Marmara region, where Europe meets Asia around the Marmara Sea, and which includes the whole of Thrace and North-west Anatolia of Turkey. The mountain, previously known as Olympus Misius, Bithynian Olympos, and Keşiş Dağı, was renamed Uludağ in 1925. The summit of Uludağ, located to the south of the city of Bursa is at 2543 m. Various vegetation types occur depending on climate type and altitude on Mt. Uludağ. As a result of the altitudinal gradient and various geological conditions, the changes from Mediterranean to Euro-Siberi-

an and alpine in the vegetation of Uludağ can clearly be seen from the bottom to the top of the mountain.

It has a Mediterranean climate with very cold winters and modified by the climatic conditions of the Black Sea region and the Inner Anatolian region. According to the data of the meteorological stations at Sarıalan (1620 m) on Uludağ, the mean annual temperature is 10 °C and mean annual rainfall is 1330 mm [14].

Populus tremula L. is a tree species in the Euro-Siberian element that grow naturally in open areas in the forest, and is distributed up to 2000 (-2300m) above sea level in Turkey [15]. It shows the distribution in open areas in the forest zone of *Abies nordmanniana* (Stev.) Spach subsp. *bornmuelleriana* (Mattf.) Coode & Cullen, *Castanea sativa* Mill., *Fagus orientalis* Lipsky from 350 to 2100 m on the northern slopes of Uludag mountain [16].

Sample collection

The study was conducted on 10 randomly selected *Populus tremula* in the location (40°06'39-42" N and 29°05'28-36"E) with an elevation of 1560-1600m at Kirazlıyayla in Uludag Mountain. Epiphytic lichen samples on the base and trunk of *Populus tremula* were collected using the methods specified by Asta et al. [17]. Lichen samples were collected from five quadrat subunits, each one has a 10x10 cm² surface area and placed on the north, south, east and west side on the base and 150 cm above the ground of the tree trunk. Sampling was performed for a total of 40 subunits per tree.

Statistical analysis

The data matrix of 40 species × 80 samples and importance values (IV) of lichens were used for statistical evaluation. The cover and frequency of occurrence of lichen species were calculated according to the north, south, east, west side on the base and trunk for individual aspen trees [18].

$$\%Frequency(F_i) = \frac{\text{the number of subunits from which species } i \text{ was recorded}}{\text{the total number of quadrat subunits examined on trees}} \times 100$$

$$\%Cover(C_i) = \frac{\text{the total cover of subunits from which species } i \text{ was recorded}}{\text{the total cover of quadrat subunits examined on trees}} \times 100$$

$$\text{Importance Value (IV)} = \%F_i + \%C_i$$

The species present at least twice on the sampling trees were evaluated statistically. A one-way analysis of variance (ANOVA) was used to test whether there is a difference in epiphytic lichen diversity on the base and trunk of *Populus tremula* and also the north, south, east and west side. Standard statistical analyses were performed using SPSS for Windows (Version 22). In all tests, the level of significance was $p \leq 0.05$. The ordination graphs according to the direction and the base-trunk of the samples were obtained with a detrended correspondence analysis (DCA). The epiphytic lichens diversity and its relationship to determined parameters (base-trunk, directions of tree trunk, circumference of tree trunk) were obtained with a canonical correspondence analysis (CCA), using the CANOCO 4.5 package [19]. The relationship between species diversity and determined parameters was determined by a Monte Carlo permutation test (495 permutations).

RESULTS

A total of 40 epiphytic lichen species were found on *Populus tremula*. Three species of them were collected only on the base of sampling trees, and ten species only on the tree trunk. Also, the 27 species were found on both the base and trunk of trees (Table 1). The most common species with the frequency of occurrence on the trees, respectively were *Lecidella elaeochroma*, *Melanohalea exasperatula*, *Caloplaca Cerina*, *Bacidia subincompta*, *Phlyctis argena*, *Physcia leptalea*, *Hypogymnia tubulosa* and *Lecanora hagenii*. The crustose growth type among the species determined is most abundant (57.5%) both the base and 150 cm above the ground on the tree trunk. The foliose growth type is 27.5% and the fruticose growth type is 15% of the determined species.

A total of 30 species were collected from the base of tree trunk. 3 species of them (*Buellia disciformis*, *Buellia Erubescens* and *Physconia enteroxantha*) were only found on the base of tree trunk. The species with the high level of importance value and the frequency of occurrence on the base of tree trunk, respectively were *Lecidella elaeochroma*, *Bacidia subincompta*, *Caloplaca Cerina*, *Phlyctis argena*, *Hypogymnia tubulosa*, *Melanohalea exasperatula*, *Lecanora hagenii* and *Rinodina exigua*. There are 23 species in the north and south of the base of tree trunk. It is to follow 17 species in the east and 20 species in the west. *Bacidia circumspecta*, *B. Subincompta*, *Caloplaca Cerina*, *Candelariella vitellina*, *Fellhanera Bouteillei*, *Hypogymnia tubulosa*, *Lecanora Chlarotera*, *L. hagenii*, *Lecidella elaeochroma*, *Melanohalea exasperatula*, *Phlyctis argena*, *Physcia leptalea* and *Rinodina exigua* are situated right in all directions on the base of trees.

A total of 37 species were collected from 150 cm above the ground on the tree trunk. 10 species of them (*Bryoria Fuscescens*, *Caloplaca holocarpa*, *Evernia prunastri*, *Micarea Melaena*, *Pertusaria Amara*, *Physcia aipolia*, *Physconia Distorta*, *Pleurosticta Acetabulum*, *Scoliosporum Chlorococcum* and *Usnea glabrescens*) were only found on the tree trunk. The species with the high level of importance value and the frequency of occurrence on the tree trunk, respectively were *Lecidella elaeochroma*, *Melanohalea exasperatula*, *Physcia leptalea*, *Caloplaca Cerina*, *Lecanora hagenii*, *Hypogymnia tubulosa*, *Phlyctis argena*, *Bacidia subincompta*, *Pseudevernia furfuracea* and *Anaptychia ciliaris*. There are 28 species in the north on 150 cm above the

ground on the tree trunk. It is to follow 26 species in the South, 21 species in the east and 27 species in the west. *Anaptychia ciliaris*, *Bacidia subincompta*, *Caloplaca Cerina*, *C. holocarpa*, *C. pollinii*, *Hypogymnia tubulosa*, *Lecania fuscella*, *Lecanora Chlarotera*, *L. hagenii*, *Lecidella elaeochroma*, *Melanohalea exasperatula*, *Phlyctis argena*, *Physcia leptalea* and *Pseudevernia furfuracea* are situated right in all directions on 150 cm above the ground on the tree trunk. The difference in epiphytic lichen diversity on the base and trunk of *Populus tremula* is not statistically significant depending on the direction.

The DCA analysis results of 80 quadrat subunits on the base and trunk of *Populus tremula* depending on the variation of importance value of 40 epiphytic lichen species are shown in Figure 1. The differences in the epiphytic lichen diversity between the base and trunk of *Populus tremula* is statistically significant, but is not different depending on the direction of the body of *Populus tremula*.

The DCA analysis results of the differences in the species diversity of the epiphytic lichens on the base and trunk of *Populus tremula* are shown in Figure 2. When compared with the species diversity between the base and trunk of *Populus tremula*, *Bacidia subincompta*, *Candelariella vitellina* and *Rinodina exigua* were statistically significant for the base. On the other hand, *Anaptychia ciliaris*, *Caloplaca holocarpa*, *Melanohalea exasperatula* and *Physcia leptalea* were significant with high importance value for the trunk. All of the fruticose species as *Anaptychia ciliaris*, *Bryoria capillaris*, *B. fuscescens*, *Evernia prunastri*, *Pseudevernia furfuracea* and *Usnea glabrescens* were found with the high importance value on the tree trunk. Generally, the foliose species on the trunk of *Populus tremula* have a higher importance values than on the base of the tree.

The two axes are represented 87.8 % of cumulative variance of species-selected parameters relation in CCA ordination (Figure 3). According to the results of Monte Carlo permutation tests, there is a significant correlation at the base-trunk in the relationship between the epiphytic lichen diversity and selected parameters. The first axis was negatively correlated with base-trunk (F:3.41, $p < 0.01$).

The species only found on the tree trunk are located on the left side of the second axis, while the species only found on the base are on the right side of the second axis of the CCA ordination plot. The species present on the base and trunk of the trees, and commonly species are located at the center of CCA ordination plot. Pearson correlation between the species and selected parameters are shown in Table 2. *Caloplaca holocarpa* and *Physcia leptalea* are on small diameter trees, while *Candelariella vitellina*, *Physconia enteroxantha* and *Rinodina exigua* are common on the large diameter trees. *Bacidia subincompta*, *Candelariella vitellina* and *Rinodina exigua* were statistically significant for the base. On the other hand, *Anaptychia ciliaris*, *Caloplaca holocarpa*, *Melanohalea exasperatula* and *Physcia leptalea* were significant for the trunk. *Bacidia subincompta*, *Candelariella vitellina* and *Rinodina exigua* at the base of *Populus tremula* have higher importance values than on the tree trunk depending on the directions. On the contrary, *Anaptychia ciliaris*, *Caloplaca holocarpa*, *Melanohalea exasperatula*, *Physcia adscendens*, *P. Leptalea* and *Scoliosporum chlorococcum* have a high importance values on the tree trunk.

DISCUSSION

The epiphytic lichen diversity on *Populus tremula* in Uludağ Mountain shows great similarity to those on *Populus tremula* in Gürgendagi Mountain which is located in the northern part of Western Anatolia [20], and in the Aladağlar National Park which is one of the largest national parks in the Mediterranean phytogeographical region of Turkey [21]. Similarly, the epiphytic lichen diversity on *Populus tremula* is very similar to previously detected on *Abies* in the same area [22].

The importance of light and humidity conditions for epiphytic vegetation is well known. At the tree level, relative humidity also tends to be higher at the trunk base and decreases with height on the tree trunk [23]. The dominance of lichens at breast height is related to different humidity levels. Except for the high humidity requires cyanolichens, lichens are more abundant in sunny environments [24]. *Caloplaca Cerina*, *C. holocarpa* and *Physcia aipolia* are characteristic indicators for xeric microclimatic conditions in sunny environments [25]. In our study, these species are found only on the tree trunk or with the high importance value on the tree trunk.

The epiphytic lichen diversity on stems and on branches was found to increase in relation to the age and diameter of the tree [26, 27]. The change in the composition of the lichen species on the tree trunks is determined by the trunk circumference on the different tree species [28, 29]. In our study, the largest differences in the epiphytic lichen diversity on *Populus tremula* were found between the trunk base and trunk.

Hypogymnia, *lecanora*, *melanelia*, *parmelia* and *physcia* are characteristic genera for sun-exposed habitats and are common genera on aspen in northern Sweden [30]. In this study, *bacidia*, *caloplaca*, *lecanora* and *physcia* are common genera on *Populus tremula*. Almost 40 species were characteristic of young aspen forest. Among them *Caloplaca Cerina*, *C. holocarpa*, *Lecanora hagenii*, *Melanelia Exasperata*, *Physcia aipolia* and *Xanthoria Parietina* [31].

Acknowledgements

This study is produced on the master thesis on the Determination of epiphytic lichen diversity on *Populus tremula* L. (European Aspen) in Uludağ Mountain (Bursa) written by Nevin HOCAOĞLU and supervisor Şaban GÜVENÇ.

REFERENCES

[1] Kivisto L, Kuusinen M. 2000. Edge effects on the epiphytic lichen flora of *picea abies* in middle boreal Finland. The Lichenologist. 32: 387-398.

[2] Rheault H, Drapeau P, Bergeron Y, Esseen PA. 2003. Edge effects on epiphytic lichen in managed black spruce forests of eastern North America. Can J For Res. 33: 23-32.

[3] Giordani P. 2006. Variables influencing the distribution of epiphytic lichens in heterogeneous areas: A case study for Liguria, NW Italy. J. Veg. Sci. 17: 195-206.

[4] Will-Wolf S, Geiser LH, Neitlich P, Reis AH. 2006. Forest lichen communities and environment – How consistent are relationships across scales? J. Veg. Sci. 17: 171-184.

[5] Hauck M. 2011. Site factors controlling epiphytic lichen abundance in northern coniferous forests. Flora. 206: 81-90.

[6] Nascimbene J, Marini L. 2010. Oak forest exploitation and black-locust invasion caused severe shifts in epi-

phytic lichen communities in Northern Italy. Science of the Total Environment. 408: 5506-5512.

[7] Nascimbene J, Nimis PL, Dainese M. 2014. Epiphytic lichen conservation in the Italian Alps: the role of forest type. Fungal Ecology. 11: 164-172.

[8] Moning C, Werth S, Dziocok F, Bassler C, Bradtka J, Hotborn T, Muller J. 2009. Lichen diversity in temperate montane forests is influenced by forest structure more than climate. For. Ecol. Manage. 258: 745-751.

[9] Spribille T, Thor G, Bunnell FL, Goward T, Bjork C. 2008. Lichens on dead wood: species–substrate relationships in the epiphytic lichen floras of the Pacific Northwest and Fennoscandia. Ecography. 31: 741-750.

[10] Liu C, Ilvesniemi H, Westman CJ. 2000. Biomass of arboreal lichens and its vertical distribution in a boreal coniferous forest in central Finland. Lichenologist. 32: 495-504.

[11] Aragon G, Martinez I, Izquierdo P, Belinchon R, Escudero A. 2010. Effects of forest management on epiphytic lichen diversity in Mediterranean forests. Applied Vegetation Science. 13: 183-194.

[12] Johansson P. 2008. Consequences of disturbance on epiphytic lichens in boreal and near boreal forests. Biol. Conserv. 141: 1933-1944.

[13] Belinchón R, Martínez I, Escudero A, Aragón G, Valladares F. 2007. Edge effects on epiphytic communities in a Mediterranean *Quercus Pyrenaica* forest. Journal of Vegetation Science. 18: 81-90.

[14] Çetin B. 1999. The moss flora of the Uludağ National Park (Bursa/Turkey). Turk J Bot. 23: 187-193.

[15] Eminağaoğlu Ö, Avcı M, Ok T, Aksoy N. 2014. *Populus L.* (Kavaklar). In: Türkiye'nin Doğal-Egzotik Ağaç ve Çalıları (ed. Akkemik Ü), s:463-470, II. Orman Genel Müdürlüğü Yayınları, Ankara.

[16] Gülyüz G. 2000. Alpine Flowers of Uludağ – Uludağ Alpin Çiçekleri. Bursa Valiliği İl Turizm Müdürlüğü, Bursa. s. 8-12.

[17] Asta J, Erhardt W, Ferretti M, Fornasier F, Kirschaum U, Nimis PL, Purvis OW, Pirintsos S, Scheidegger C, van Haluwyn C, Wirth V. 2002. Mapping lichen diversity as an indicator of environmental quality. In: Monitoring with Lichens- Monitoring Lichens. (ed. Nimis PL, Scheidegger C, Wolseley PA), Nato Science Program-IV, Vol. VII, pp. 273-279, Kluwer Academic Publisher, The Netherlands.

[18] Ellis CJ, Coppins BJ. 2006. Contrasting functional traits maintain lichen epiphyte diversity in response to climate and autogenic succession. Journal of Biogeography. 33: 1643-1656.

[19] Ter Braak CJF. 1995. Ordination. In: Data Analysis in Community and Landscape Ecology (ed. Jongman RHG, Ter Braak CJF, Van Tongeren OFR), pp. 91-173, Cambridge University Press, Cambridge, UK.

[20] Çobanoğlu G, Sevgi O. 2006. Contribution to the Lichen Flora of Gürgen Dağı (Çanakkale). Turk J Bot. 30: 47-54.

[21] Halıcı MG, Aksoy A. 2009. Lichenised and Lichenicolous Fungi of Aladağlar National Park (Niğde, Kayseri and Adana Provinces) in Turkey. Turk J Bot. 33: 169-189.

[22] Öztürk Ş, Güvenç Ş. 2010. The distribution of epiphytic lichens on Uludağ fir (*Abies nordmanniana* (Steven) Spach subsp. *Bornmuelleriana* (Mattf.) Coode & Cullen) forests along an altitudinal gradient (Mt. Uludağ, Bursa, Turkey). Ekoloji. 19(74): 131-138.

[23] Barkman JJ. 1958. Phytosociology and ecology of

Cryptogamic epiphytes. Van Gorcum & comp NV., Assen, Netherlands.

[24] Gustafsson L, Eriksson I. 1995. Factors of Importance for the Epiphytic Vegetation of Aspen *Populus tremula* with Special Emphasis on Bark Chemistry and Soil Chemistry. *Journal of Applied Ecology*. 32(2): 412-424.

[25] Boudreault C, Coxson DS, Vincent E, Bergeron Y, Marsh J. 2008. Variation in epiphytic lichen and bryophyte composition and diversity along a gradient of productivity in *Populus tremuloides* stands of northeastern British Columbia, Canada. *Ecoscience*. 15(1): 101-112.

[26] Hedenas H, Ericson L. 2000. Epiphytic macrolichens as conservation indicators: successional sequence in *Populus tremula* stands. *Biological Conservation*. 93: 43-53.

[27] Lie MH, Arup U, Grytnes JA, Ohlson M. 2009. The importance of host tree age, size and growth rate as determinants of epiphytic lichen diversity in boreal spruce forests. *Biodivers Conserv*. 18: 3579-3596

[28] Jürriado I, Liira J, Paal J. 2009. Diversity of epiphytic lichens in boreo-nemoral forests on the North-Estonian limestone escarpment: the effect of tree level factors and

local environmental conditions. *The Lichenologist*. 41(1): 81-96.

[29] Thor G, Johansson P, Jönsson MT. 2010. Lichen diversity and red-listed lichen species relationships with tree species and diameter in wooded meadows. *Biodivers Conserv*. 19: 2307-2328.

[30] Hedenas H, Ericson L. 2004. Aspen lichens in agricultural and forest landscapes: the importance of habitat quality. *Ecography*. 27: 1-11.

[31] Lundström J, Jonsson F, Perhans K, Gustafsson L. 2013. Lichen species richness on retained aspens increases with time since clear-cutting. *Forest Ecology and Management*. 293: 49-56.

Table 1. Mean ± Standard deviation (SD) of the importance value (IV) and frequency (F) of occurrence of epiphytic lichen species on the base and trunk of *Populus tremula*. The results were obtained by one-way ANOVA (n=40, df=1)

Sampling location of tree trunk	Abbreviations of species name	BASE										TRUNK										One-Way	
		154±6,70										138±4,94											
Mean±SD of circumference of tree trunk		North		South		East		West		IV	North		South		East		West		IV	ANOVA			
Direction of tree trunk		F	IV	F	IV	F	IV	F	IV	Mean±SD	F	IV	F	IV	F	IV	F	IV	Mean±SD	F	Sig.		
Species																							
Anaptychia ciliaris	Anap cil	2	4,1	1	2,1					1,6±5,5	5	10,7	2	4,3	5	10,3	9	18,6	10,9±21,6	7,115	,009		
Bacidia arceutina	Baci arc			3	6,0	1	2,0	1	2,0	2,5±8,1	1	2,0							0,5±3,2	2,131	,148		
Bacidia circumspecta	Baci cir	2	4,0	1	2,0	1	2,0	1	2,0	2,5±6,7	1	2,0					2	4,0	1,5±5,4	0,552	,460		
Bacidia subincompta	Baci sub	10	23,8	16	34,3	19	45,3	5	10,1	28,4±30,7	6	12,2	7	14,2	7	14,2	5	10,1	12,7±24,6	6,366	,014		
Bryoria capillaris	Byro cap	2	4,1							1,0±4,5	6	13,7					1	2,2	3,9±18,8	0,930	,338		
Bryoria fuscescens	Byro fus										3	6,5	2	5,1			1	2,3	3,5±13,2	2,738	,102		
Buellia disciformis	Buel dis	2	4,1			4	8,2			3,1±10,9										3,162	,079		
Buellia erubescens	Buel eru	1	2,0	2	4,0					1,5±7,1										1,836	,179		
Caloplaca cerina	Calo cer	7	14,1	11	22,2	8	16,1	13	26,2	19,6±23,9	11	22,1	15	30,2	7	14,1	14	28,2	23,6±26,9	0,496	,483		
Caloplaca herbidella	Calo her			1	2,0					0,5±3,2			1	2,1					0,5±3,3	0,001	,981		
Caloplaca holocarpa	Calo hol										6	12,1	6	12,1	2	4,0	3	6,0	8,5±16,9	10,148	,002		
Caloplaca pollinii	Calo pol	1	2,0	4	8,1			2	4,0	3,5±12,0	2	4,0	5	10,1	5	10,1	2	4,0	7,1±14,8	1,370	,245		
Candelariella vitellina	Cand vit	4	8,1	2	4,0	5	10,1	4	8,1	7,6±16,9	1	2,0							0,5±3,2	6,738	,011		
Evernia prunastri	Ever pru												1	2,1	2	4,3			1,6±7,4	1,829	,180		
Fellhanera bouteillei	Fell bou	1	2,0	3	6,1	4	8,1	3	6,1	5,6±12,1	1	2,0	3	6,1			1	2,0	2,5±8,2	1,708	,195		
Hypogymnia tubulosa	Hypo tub	8	17,2	9	18,4	3	6,1	7	14,2	13,9±27,6	15	31,5	9	18,8	7	14,7	9	18,5	20,8±35,5	0,939	,336		
Lecania fuscella	Leca fus			1	2,0	2	4,0	3	6,0	3,0±10,7	3	6,0	7	14,2	3	6,1	4	8,0	8,6±15,7	3,396	,069		
Lecanora carpinea	Leca car	1	2,0					3	6,2	2,1±7,7	2	4,0					1	2,0	1,5±5,4	0,134	,715		
Lecanora chlorotera	Leca chl	3	6,2	5	10,3	4	8,3	2	4,2	7,3±16,6	4	8,2	2	4,3	5	10,2	1	2,0	6,2±13,3	0,101	,752		

Lecanora hagenii	Leca hag	3	6,0	10	20,2	6	12,2	5	10,1	12,1±22,3	7	14,2	17	34,4	11	22,3	8	16,2	21,7±34,7	2,184	,143
Lecanora symmicta	Leca sym	4	8,3	1	2,1					2,6±13,5			1	2,0					0,5±3,2	0,925	,339
Lecidella elaeo-chroma	Leci ela	14	28,9	22	44,7	20	40,6	14	28,5	35,7±31,4	19	38,4	24	49,0	19	38,5	20	40,7	42,7±29,2	1,070	,304
Melanelixia glabrata	Mela gla	3	6,1							1,5±7,2	1	2,0					1	2,3	1,1±4,8	0,109	,742
Melanohalea exasperatula	Melo exa	8	16,3	5	10,1	5	10,1	7	14,2	12,6±20,8	19	39,1	17	35,4	16	32,8	20	40,6	36,9±36,6	13,308	,000
Micarea melaena	Mica mel										3	6,1			1	2,0	3	6,0	3,5±13,6	2,694	,105
Parmelia sulcata	Parm sul					1	2,1			0,5±3,3	1	2,1					1	2,0	1,0±4,6	0,339	,562
Pertusaria amara	Pert ama										2	4,1					1	2,0	1,5±7,1	1,834	,180
Phlyctis argena	Phly arg	8	17,1	13	27,8	10	21,2	8	16,6	20,6±32,6	7	14,6	8	16,7	9	18,9	7	14,6	16,2±24,2	0,482	,489
Physcia adscendens	Phys ads							1	2,1	0,5±3,3			2	4,0	1	2,0	3	6,1	3,0±9,7	2,393	,126
Physcia aipolia	Phys aip												3	6,2	2	4,1			2,6±10,7	2,326	,131
Physcia leptalea	Phys lep	5	10,1	2	4,0	4	8,0	2	4,1	6,6±12,4	15	30,3	16	32,3	9	18,2	15	30,5	27,8±31,3	15,927	,000
Physconia distorta	Pyco dis												1	2,2			1	2,0	1,1±4,7	2,047	,157
Physconia enteroxantha	Pyco ent	1	2,0	1	2,0					1,0±4,5										2,053	,156
Pleurosticta acetabulum	Pleu ace										4	9,3	2	4,3					3,4±16,0	1,792	,185
Pseudevernia furfuracea	Pseu fur	5	10,4	3	6,1			1	2,0	4,6±17,8	5	12,6	8	16,4	4	8,8	6	12,8	12,6±31,2	1,993	,162
Rinodina capensis	Rino cap							2	4,0	1,0±6,4	2	4,0			1	2,0	1	2,0	2,0±7,6	0,412	,523
Rinodina exigua	Rino exi	4	8,1	6	12,1	10	20,1	1	2,0	10,6±22,8			1	2,0					0,5±3,2	7,661	,007
Scoliciosporum chlorococcum	Scol chl														3	6,1	1	2,0	2,0±7,6	2,793	,099

Table 2. Pearson correlation coefficients between the species and selected parameters.

Species	Selected parameters		
	Circumference of tree trunk	Base-Trunk	Direction of tree trunk
<i>Anaptychia ciliaris</i>	-0.185	0.289**	0.278*
<i>Bacidia subincompta</i>	0.180	-0.275*	-0.275*
<i>Caloplaca holocarpa</i>	-0.269*	0.339**	0.239*
<i>Candelariella vitellina</i>	0.397**	-0.282*	-0.46*
<i>Melanohalea exasperatula</i>	-0.210	0.382**	0.330**
<i>Physcia adscendens</i>	-0.141	0.173	0.234*
<i>Physcia leptalea</i>	-0.259*	0.412**	0.330**
<i>Physconia enteroxantha</i>	0.338**	-0.160	-0.209
<i>Rinodina exigua</i>	0.293**	-0.299**	-0.281*
<i>Scoliciosporum chlorococcum</i>	-0.147	0.186	0.223*

Correlation is significant at the 0.01 level (**) and at the 0.05 level (*)

Figure 1. DCA ordination of 80 quadrat subunits on the base and trunk of *Populus tremula* depending on the variation of importance value of 40 epiphytic lichen species. Total inertia in species data: 4,40. Eigenvalues: 0.59 (axis 1), 0.38 (axis 2). Length of gradient: 4.99 (axis 1), 3.70 (axis 2).



